




enbond



Tissue Adhesive



A Revolution in wound closure

- Quick and easy
- Less scarring
- Anti-septic action
- Safe and Precise

A Cosmetic Revolution in Surgical procedures

The procedure ends just when it starts

Easy, Simple and Effective



www.nectarlabs.in

Business Queries
exports@nectarlabs.in

Manufactured in India by:



Nectar Laboratories Private Limited

3rd Floor, TGV Mansion, 6-2-1012, Khairatabad, Hyderabad - 500 004, Telangana, India.

For Easy, Simple and Effective Wound Closure

Use **enbond**
(n-Butyl 2-Cyanoacrylate)

A Cosmetic revolution in Surgical procedures



For Easy, Simple and Effective Wound Closure

Use **enbond**
(n-Butyl 2-Cyanoacrylate)

A Cosmetic revolution in Surgical procedures



THE PRODUCT

enbond is n - Butyl 2 - Cyanoacrylate and is a medical grade Bio-adhesive.

Cyanoacrylates are derivatives of acrylic acid. It is liquid in its monomer form and polymerises quickly as it comes into contact with water or body moisture.

The polymeric matter forms a thin layer on the skin which holds the skin edges together tightly.

This Property enables the use of **enbond** in the closure of incisional as well as post-traumatic wounds.

Presentation:
enbonds available in ampoules of 0.25 ml / 0.5 ml / 1 ml.



BENEFITS

BIO-COMPATIBLE: Butyl cyanoacrylates are inert and non-toxic. They produce less oedema, inflammation and granulation than sutures.

QUICK ADHESION: Binds tissues faster than sutures. It polymerises & solidifies within a minute on contact with body moisture and seals the wound almost immediately.

EXCELLENT HAEMOSTATIC EFFECT: When injected endoscopically into the bleeding varices is reported to stop the bleeding in upto 93% cases within a single session.

ENHANCED COSMETIC APPEAL: Single linear scar was seen after (n-Butyl 2 Cyanoacrylate) use.



How
to
Use

APPLICATIONS

Closure of external surgical, post-traumatic wounds, Cuts and Lacerations

Closure of External incisional wounds in







Plastic and Cosmetic Surgery

enbond is widely used in:

Gynaecology & obstetrics	All facial Surgeries
Appendectomy	Surgeries to remove scars
Cholecystectomy	Correction of ear lobe punctures
G.I Resection Biopsy	Corrective breast surgery
Gastro-Jejunostomy	
Other abdominal surgeries, Laproscopy incisions etc.	

1. Remove enbond package from storage and bring to room temperature. Remove the ampoule from the blister.
2. Clean the oozing blood and dry the wound area as fast as possible.
3. Take the enbond ampoule and snap the tip at the marked area holding it vertically with both the hands.
4. Approximate the wound edges precisely. Forceps can be used to hold the edges of the skin in case of larger wounds.
5. Apply a thin layer of enbond throughout the approximated wound and allow it to dry for about 2 minutes.

DO'S & DON'TS



Enbond must be used only when in liquid form. The condition of enbond can be assessed before opening the adhesive ampoule. If found viscous, discard the product.